Stockholm 2018

Contribution ID: 9

Type: Contributed poster

Time Dynamics and Spectral Resolved Emission Imaging of Colliding Laser Produced Aluminum Plasma

Monday, 25 June 2018 18:00 (15 minutes)

Time Dynamics and Spectral Resolved Emission Imaging of Colliding Laser Produced Aluminum Plasma Haider Mahdi Al-Juboori1, Thomas McCormack1 11 Iniversity College Dublin, School of Physics, Bolfold, Dublin 4, Iroland

1University College Dublin, School of Physics, Belfield, Dublin 4, Ireland. email: haider.al-juboori@ucd.ie

Abstract

Colliding laser produced plasma have been investigated in a wide range of laser irradiance regimes (1010- 1015 W/cm2) and for several purposes: by employing them as possible fuels for ICF-inertial fusion confinement or in order to reproduce, on a laboratory-scale, astronomical processes as for example collision-less shock eaves production [1,2].

In the present work, the collision of two aluminum plasmas was investigated by combining both time and space resolved spectroscopy.

Plasma plumes were produced by a Continuum TM Surelite Nd:YAG laser system with pulse duration of FWHM of 6 ns and wavelength of 1064 nm, at a laser irradiance of ~1011 W/cm3 on slab Al target.

By using Filter sensitive technique, the temporally and spatially resolved electron density and temperature at the stagnation layer were present, with a time resolution of 10ns.

The data analysis confirms that the electron density of the stagnation layer evolves over a longer timescale than in the single plume case.

References:

[1] A.S. Wan, T.W. Barbee, R. Cauble, P. Celliers, L.B.D. Silva, J.C. Moreno, P.W. Rambo, G.F. Stone, J.E. Trebes, F. Weber, Physical Review E 55 (1997) 6293–6296.

[2] N.C. Woolsey, C. Courtois, R.O. Dendy, Plasma Physics and Controlled Fusion 46 (2004) B397.

Primary authors: Mr AL-JUBOORI, Haider (UCD); Dr MCCORMACK, Thomas (UCD)

Presenter: Mr AL-JUBOORI, Haider (UCD)

Session Classification: Poster session