



Contribution ID: 17

Type: Invited oral

## Molecules at high x-ray intensity: Challenges for theory

Monday, 25 June 2018 15:00 (30 minutes)

One of the key opportunities offered by the development of x-ray free-electron lasers is the determination, at atomic resolution, of the three-dimensional structure of biologically relevant macromolecules. The basic idea underlying molecular imaging using x-ray free-electron lasers is the “diffract-and-destroy” concept: If one uses an x-ray pulse that is sufficiently short (on the order of femtoseconds), then in a single shot an x-ray scattering pattern may be obtained that is practically unaffected by atomic displacements triggered by ionization events during the x-ray pulse. What cannot be eliminated in this way is the impact of the electronic damage on the x-ray scattering patterns. Theory, therefore, plays an important role in the development of this new imaging technique: A quantitative understanding is required of the damage processes occurring during the exposure of a molecule to an ultraintense, ultrafast x-ray pulse. In this talk, I will present progress we have made in order to address this challenge. One tool we have developed, XMDYN [1], is a molecular-dynamics code that utilizes ab-initio atomic electronic-structure information, computed on the fly, within a Monte-Carlo framework. XMDYN has been successfully tested through experiments at LCLS [2] and SACLA [3]. XMDYN is part of a powerful start-to-end simulation framework for single-particle imaging at the European XFEL [4,5]. Recently, we have taken first steps towards a full ab-initio framework for simulating high-intensity x-ray/matter interactions [6,7]. Our new XMOLECULE software solves the polyatomic quantum-mechanical electronic-structure problem for every electronic state arising during the exposure of a molecule to a strong x-ray pulse. From this information, electronic transition rates (such as Auger decay rates) are computed on the fly, and the associated rate equations are integrated utilizing a Monte-Carlo method. XMOLECULE played a key role in a recent LCLS experiment on iodomethane, in which hard x-rays focused to a peak intensity exceeding  $10^{19}$  W/cm<sup>2</sup> produced the highest charge states ever formed using light [8]. Not only did XMOLECULE correctly predict the charge-state distribution observed, but it also helped identify a new molecular ionization enhancement mechanism based on intramolecular charge transfer.

- [1] Z. Jurek *et al.*, *J. Appl. Cryst.* **49**, 1048 (2016).
- [2] B. F. Murphy *et al.*, *Nature Commun.* **5**, 4281 (2014).
- [3] T. Tachibana *et al.*, *Sci. Rep.* **5**, 10977 (2015).
- [4] C. H. Yoon *et al.*, *Sci. Rep.* **6**, 24791 (2016).
- [5] C. Fortmann-Grote *et al.*, *IUCr* **4**, 560 (2017).
- [6] Y. Hao *et al.*, *Struct. Dyn.* **2**, 041707 (2015).
- [7] L. Inhester *et al.*, *Phys. Rev. A* **94**, 023422 (2016).
- [8] A. Rudenko *et al.*, *Nature* **546**, 129 (2017).

**Primary author:** SANTRA, Robin

**Presenter:** SANTRA, Robin

**Session Classification:** New developments